

BUMBLE BEES

APPEARANCE

The British Isles are host to 19 species of bumble bee. These vary in size and colour with the most common being the Red Tailed, White Tailed and Buff Tailed bumble bees. Size in each species can vary dramatically from a worker to the large queen. All species are beneficial as pollinators while collecting nectar and pollen from their chosen food plant, some of which are solely pollinated by bumble bees.



LIFE CYCLE

Queens emerge in early spring when a suitable nest chamber is found (an old mouse nest in a sheltered place or an old nest in a bird box). The young queen builds a small pollen chamber where the first eggs are laid and these develop into workers. By mid-summer the colony may contain between 50 - 200 workers. Later eggs develop into males and queens and these males and the workers eventually die leaving only queens to hibernate over winter to start a new colony in the following spring.

FEEDING HABITS

Queens emerging in spring depend on early flowering plants such as white dead nettle, flowering currant, willows, and pussy willow. Tongue length in the different species varies tremendously enabling some species to feed on plants which are inaccessible to any other bees.

SIGNS OF INFESTATION

After a queen has colonised a chosen site continual movement to and from the nest site can be observed.

For further information please contact:

- 📞 Pest Control
Commercial Services
Nottingham City Council
Eastcroft Depot
- ☎ 0115 915 2020
- ✉ pest.control@nottinghamcity.gov.uk

Or visit our Commercial Services website at:

www.nottmcommercialservices.co.uk



Nottingham
City Council

Pest
Control